


Doctor Discussion Guide

Print this guide to fill out and bring to your next doctor's appointment. Your answers could help you have a more informed conversation about your weight and prescription treatment options.




Questions to Ask Your Doctor

- Is a prescription weight-loss option right for me?
- How does BELVIQ work?
- What results can I expect with BELVIQ?
- What are the side effects of BELVIQ?

What's Your BMI?

Your body mass index, or BMI, is a number that can help you determine if you're within a normal weight range. Use the chart below to find your BMI. If it falls within a shaded blue or red area, ask your doctor if BELVIQ may be right for you.



What is BELVIQ[®]?

BELVIQ is an FDA-approved prescription weight-loss medication that, when used with diet and exercise, can help some overweight* adults with a weight-related medical problem, or obese adults, lose weight and keep it off.

It is not known if BELVIQ when taken with other prescription, over-the-counter, or herbal weight-loss products is safe and effective. It is not known if BELVIQ changes your risk of heart problems, stroke, or death due to heart problems or stroke.

*Overweight (body mass index [BMI] of 27 kg/m² or greater) with at least one weight-related medical condition, such as high blood pressure, high cholesterol, or type 2 diabetes; Obese (BMI of 30 kg/m² or greater).

Important Safety Information

- **Pregnancy:** Do not take BELVIQ if you are pregnant or planning to become pregnant, as weight loss offers no potential benefit during pregnancy and BELVIQ may harm your unborn baby.

Please see additional Important Safety Information on the following pages. For more information about BELVIQ[®], talk to your doctor and see available Product Information.

Information to Give Your Doctor

For each question, check all of the boxes that apply to you. Then bring this completed sheet to your next doctor's appointment to discuss your answers.

Do you have any of the following weight-related medical conditions?

- High blood pressure
- High cholesterol
- Type 2 diabetes
- Other _____

Do you have a family history of any of the following medical conditions?

- High blood pressure
- High cholesterol
- Type 2 diabetes
- Other _____

What have you done to try to lose weight in the past?

- Diet or weight-loss program
- Over-the-counter weight-loss pills
- Weight-loss supplements
- Meal replacement programs
- I've never tried to lose weight

How would you describe your past attempts at weight loss?

- Successful
- Unsuccessful
- I've lost weight in the past, but have gained it back
- I've never tried to lose weight

Are you currently on any kind of diet?

- I am on a diet or weight-loss program
- I take over-the-counter weight-loss pills or supplements
- I use a meal replacement program
- I count calories
- I am not currently dieting

What is your current level of physical activity?

- I exercise every day
- I exercise a few times a week
- I rarely exercise
- I never exercise

- I am committed to losing weight through diet and exercise and exploring prescription treatment options.**

Important Safety Information Continued

- **Serotonin Syndrome or Neuroleptic Malignant Syndrome (NMS)-like reactions:** Before using BELVIQ, tell your doctor about all the medicines you take, especially medicines that treat depression, migraines, mental problems, or the common cold. These medicines may cause serious or life-threatening side effects if taken with BELVIQ. Call your doctor right away if you experience agitation, hallucinations, confusion, or other changes in mental status; coordination problems; uncontrolled muscle spasms; muscle twitching; restlessness; racing or fast heartbeat; high or low blood pressure; sweating; fever; nausea; vomiting; diarrhea; or stiff muscles.

Important Safety Information Continued

- **Valvular heart disease:** Some people taking medicines like BELVIQ have had heart valve problems. Call your doctor right away if you experience trouble breathing; swelling of the arms, legs, ankles, or feet; dizziness, fatigue, or weakness that will not go away; or fast or irregular heartbeat. Before taking BELVIQ, tell your doctor if you have or have had heart problems.
- **Changes in attention or memory:** BELVIQ may slow your thinking. You should not drive a car or operate heavy equipment until you know how BELVIQ affects you.
- **Mental problems:** Taking too much BELVIQ may cause hallucinations, a feeling of being high or in a very good mood, or feelings of standing outside your body.
- **Depression or thoughts of suicide:** Call your doctor right away if you notice any mental changes, especially sudden changes in your mood, behaviors, thoughts, or feelings, or if you have depression or thoughts of suicide.
- **Low blood sugar:** Weight loss can cause low blood sugar in people taking medicines for type 2 diabetes, such as insulin or sulfonylureas. Blood sugar levels should be checked before and while taking BELVIQ. Changes to diabetes medication may be needed if low blood sugar develops.
- **Painful erections:** If you have an erection lasting more than 4 hours while on BELVIQ, stop taking BELVIQ and call your doctor or go to the nearest emergency room right away.
- **Slow heartbeat:** BELVIQ may cause your heart to beat slower.
- **Decreases in blood cell count:** BELVIQ may cause your red and white blood cell counts to decrease.
- **Increase in prolactin:** BELVIQ may increase the amount of a hormone called prolactin. Tell your doctor if your breasts begin to make milk or a milky fluid, or if you are a male and your breasts increase in size.
- **Most common side effects in patients without diabetes:** Headache, dizziness, fatigue, nausea, dry mouth, and constipation.
- **Most common side effects in patients with diabetes:** Low blood sugar, headache, back pain, cough, and fatigue.
- **Nursing:** BELVIQ should not be taken while breastfeeding.
- **Drug interactions:** Before taking BELVIQ, tell your doctor if you take medicines for depression, migraines, or other medical conditions, such as: triptans; medicines used to treat mood, anxiety, psychotic or thought disorders, including tricyclics, lithium, selective serotonin reuptake inhibitors, selective serotonin-norepinephrine reuptake inhibitors, monoamine oxidase inhibitors, or antipsychotics; cabergoline; linezolid (an antibiotic); tramadol; dextromethorphan (an over-the-counter (OTC) common cold/cough medicine); OTC supplements such as tryptophan or St. John's Wort; or erectile dysfunction medicines.

BELVIQ is a federally controlled substance (CIV) because it may be abused or lead to drug dependence.

For more information about BELVIQ®, talk to your doctor and see attached Product Information, or visit the BELVIQ website for full [Product Information](#).

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

IMPORTANT PATIENT INFORMATION

Read the Patient Information that comes with BELVIQ® (BEL-VEEK) (lorcaserin hydrochloride) tablets before you start taking it and each time you get a refill. There may be new information. This page does not take the place of talking with your doctor about your medical condition or treatment. If you have any questions about BELVIQ, talk to your doctor or pharmacist.

What is BELVIQ?

BELVIQ is a prescription medicine that may help some obese adults or overweight adults who also have weight related medical problems lose weight and keep the weight off. BELVIQ should be used with a reduced calorie diet and increased physical activity.

It is not known if BELVIQ is safe and effective when taken with other prescription, over-the-counter, or herbal weight loss products.

It is not known if BELVIQ changes your risk of heart problems or stroke or of death due to heart problems or stroke.

It is not known if BELVIQ is safe when taken with some other medicines that treat depression, migraines, mental problems, or the common cold (serotonergic or antidopaminergic agents).

It is not known if BELVIQ is safe and effective in children under 18 years old.

BELVIQ is a federally controlled substance (CIV) because it contains lorcaserin hydrochloride and may be abused or lead to drug dependence. Keep your BELVIQ in a safe place, to protect it from theft. Never give your BELVIQ to anyone else, because it may cause harm to them. Selling or giving away this medicine is against the law.

Who should not take BELVIQ?

Do not take BELVIQ if you:

- are pregnant or planning to become pregnant. BELVIQ may harm your unborn baby.

What should I tell my healthcare provider before taking BELVIQ?

Before you take BELVIQ, tell your doctor if you:

- **have or have had heart problems including:**
 - congestive heart failure
 - heart valve problems
 - slow heartbeat or heart block
- have diabetes
- have a condition such as sickle cell anemia, multiple myeloma, or leukemia
- have a deformed penis, Peyronie's disease, or ever had an erection that lasted more than 4 hours
- have kidney problems
- have liver problems
- are pregnant or plan to become pregnant
- are breastfeeding or plan to breastfeed. It is not known if BELVIQ passes into your breastmilk. You and your doctor should decide if you will take BELVIQ or breastfeed. You should not do both.

Tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements.

BELVIQ may affect the way other medicines work, and other medicines may affect how BELVIQ works.

Especially tell your doctor if you take medicines for depression, migraines or other medical conditions such as:

- triptans, used to treat migraine headache
- medicines used to treat mood, anxiety, psychotic or thought disorders, including tricyclics, lithium, selective serotonin reuptake inhibitors (SSRIs), selective serotonin-norepinephrine reuptake inhibitors (SNRIs), monoamine oxidase inhibitors (MAOIs), or antipsychotics
- cabergoline
- linezolid, an antibiotic
- tramadol
- dextromethorphan, an over-the-counter medicine used to treat the common cold or cough

- over-the-counter supplements such as tryptophan or St. John's Wort
- medicines to treat erectile dysfunction

Ask your doctor or pharmacist for a list of these medicines, if you are not sure.

Know all the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

How should I take BELVIQ?

- Take BELVIQ exactly as your doctor tells you to take it.
- Your doctor will tell you how much BELVIQ to take and when to take it.
 - Take 1 tablet 2 times each day.
 - **Do not** increase your dose of BELVIQ.
 - BELVIQ can be taken with or without food.
- Your doctor should start you on a diet and exercise program when you start taking BELVIQ. Stay on this program while you are taking BELVIQ.
- Your doctor should tell you to stop taking BELVIQ if you do not lose a certain amount of weight within the first 12 weeks of treatment.
- If you take too much BELVIQ or overdose, call your doctor or go to the nearest emergency room right away.

What should I avoid while taking BELVIQ?

- **Do not** drive a car or operate heavy machinery until you know how BELVIQ affects you. BELVIQ can slow your thinking.

What are the possible side effects of BELVIQ?

BELVIQ may cause serious side effects, including:

- **Serotonin Syndrome or Neuroleptic Malignant Syndrome (NMS)-like reactions.** BELVIQ and certain medicines for depression, migraine, the common cold, or other medical problems may affect each other causing serious or life-threatening side effects. Call your doctor right away if you start to have any of the following symptoms while taking BELVIQ:
 - mental changes such as agitation, hallucinations, confusion, or other changes in mental status
 - coordination problems, uncontrolled muscle spasms, or muscle twitching (overactive reflexes)
 - restlessness
 - racing or fast heartbeat, high or low blood pressure
 - sweating or fever
 - nausea, vomiting, or diarrhea
 - muscle rigidity (stiff muscles)
- **Valvular heart disease.** Some people taking medicines like BELVIQ have had problems with the valves in their heart. Call your doctor right away if you have any of the following symptoms while taking BELVIQ:
 - trouble breathing
 - swelling of the arms, legs, ankles, or feet
 - dizziness, fatigue, or weakness that will not go away
 - fast or irregular heartbeat
- **Changes in your attention or memory.**
- **Mental problems.** Taking BELVIQ in high doses may cause psychiatric problems such as:
 - hallucinations
 - feeling high or in a very good mood (euphoria)
 - feelings of standing next to yourself or out of your body (disassociation)
- **Depression or thoughts of suicide.** You should pay attention to any mental changes, especially sudden changes, in your mood, behaviors, thoughts, or feelings. Call your healthcare provider right away if you have any mental changes that are new, worse, or worry you.
- **Low blood sugar (hypoglycemia) in people with type 2 diabetes mellitus who also take medicines used to treat type 2 diabetes mellitus.** Weight loss can cause low blood sugar in people with type 2 diabetes mellitus who also take medicines used to treat type 2 diabetes mellitus (such as insulin or sulfonylureas). You should check your blood sugar before you start taking BELVIQ and while you take BELVIQ.
- **Painful erections (priapism).** The medicine in BELVIQ

can cause painful erections that last more than 6 hours. If you have an erection lasting more than 4 hours whether it is painful or not, stop using BELVIQ and call your doctor or go to the nearest emergency room right away.

- **Slow heartbeat.** BELVIQ may cause your heart to beat slower. Tell your doctor if you have a history of your heart beating slow or heart block.
- **Decreases in your blood cell count.** BELVIQ may cause your red and white blood cell count to decrease. Your doctor may do tests to check your blood cell count while you are taking BELVIQ.
- **Increase in prolactin.** The medicine in BELVIQ may increase the amount of a certain hormone your body makes called prolactin. Tell your doctor if your breasts begin to make milk or a milky discharge or if you are a male and your breasts begin to increase in size.

The most common side effects of BELVIQ include:

- headache
- dizziness
- fatigue
- nausea
- dry mouth
- constipation
- cough
- low blood sugar (hypoglycemia) in patients with diabetes
- back pain

Tell your doctor if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of BELVIQ. For more information, ask your doctor or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How do I store BELVIQ?

Store BELVIQ at room temperature between 59°F to 86°F (15°C to 30°C).

Safely throw away medicine that is out of date or no longer needed.

Keep BELVIQ and all medicines out of the reach of children.

General information about the safe and effective use of BELVIQ.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use BELVIQ for a condition for which it was not prescribed. Do not give BELVIQ to other people, even if they have the same symptoms you have. It may harm them.

This Patient Information summarizes the most important information about BELVIQ. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about BELVIQ that is written for health professionals.

For more information, go to www.BELVIQ.com Website or call 1-888-274-2378.

What are the ingredients in BELVIQ?

Active Ingredient: lorcaserin hydrochloride

Inactive Ingredients: silicified microcrystalline cellulose; hydroxypropyl cellulose NF; croscarmellose sodium NF; colloidal silicon dioxide NF; polyvinyl alcohol USP; polyethylene glycol NF; titanium dioxide USP; talc USP; FD&C Blue #2 aluminum lake; and magnesium stearate NF.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Rx Only

BELVIQ® is a registered trademark of Arena Pharmaceuticals GmbH, Zofingen, Switzerland
Manufactured by Arena Pharmaceuticals GmbH, Untere Brühlstrasse 4, CH-4800, Zofingen, Switzerland
Distributed by Eisai Inc., Woodcliff Lake, NJ 07677

©2013 Eisai Inc. BELV1138 11/13